

WHAT IS A DEPRESSIVE DISORDER?

A depressive disorder is an illness that involves the body, mood, and thoughts. It affects the way a person eats and sleeps, the way one feels about oneself, and the way one thinks about things. A depressive disorder is not the same as a passing blue mood. It is not a sign of personal weakness or a condition that can be willed or wished away. People with a depressive illness cannot merely "pull themselves together" and get better. Without treatment, symptoms can last for weeks, months, or years. Appropriate treatment, however, can help most people who suffer from depression.

TYPES OF DEPRESSION

Depressive disorders come in different forms, just as is the case with other illnesses such as heart disease. This pamphlet briefly describes three of the most common types of depressive disorders. However, within these types there are variations in the number of symptoms, their severity, and persistence.

Major depression is manifested by a combination of symptoms (see symptom list) that interfere with the ability to work, study, sleep, eat, and enjoy once pleasurable activities. Such a disabling episode of depression may occur only once but more commonly occurs several times in a lifetime.

A less severe type of depression, dysthymia, involves long-term, chronic symptoms that do not disable, but keep one from functioning well or from feeling good. Many people with dysthymia also experience major depressive episodes at some time in their lives.

Another type of depression is bipolar disorder, also called manic-depressive illness. Not nearly as prevalent as other forms of depressive disorders, bipolar disorder is characterized by cycling mood changes: severe highs (mania) and lows (depression). Sometimes the mood switches are dramatic and rapid, but most often they are gradual. When in the depressed cycle, an individual can have any or all of the symptoms of a depressive disorder. When in the manic cycle, the individual may be overactive, overtalkative, and have a great deal of energy. Mania often affects thinking, judgment, and social behavior in ways that cause serious problems and embarrassment. For example, the individual in a manic phase may feel elated, full of grand schemes that might range from unwise business decisions to romantic sprees. Mania, left untreated, may worsen to a psychotic state.

SYMPTOMS OF DEPRESSION AND MANIA

Not everyone who is depressed or manic experiences every symptom. Some people experience a few symptoms, some many. Severity of symptoms varies with individuals and also varies over time.

Depression

Persistent sad, anxious, or "empty" mood
Feelings of hopelessness, pessimism
Feelings of guilt, worthlessness, helplessness
Loss of interest or pleasure in hobbies and activities that were once enjoyed, including sex
Decreased energy, fatigue, being "slowed down"
Difficulty concentrating, remembering, making decisions
Insomnia, early-morning awakening, or oversleeping
Appetite and/or weight loss or overeating and weight gain
Thoughts of death or suicide; suicide attempts
Restlessness, irritability
Persistent physical symptoms that do not respond to treatment, such as headaches, digestive disorders, and chronic pain

Mania

Abnormal or excessive elation
Unusual irritability
Decreased need for sleep
Grandiose notions
Increased talking
Racing thoughts
Increased sexual desire
Markedly increased energy
Poor judgment
Inappropriate social behavior

Depression in Women

Women experience depression about twice as often as men. Many hormonal factors may contribute to the increased rate of depression in women-particularly such factors as menstrual cycle changes, pregnancy, miscarriage, postpartum period, pre-menopause, and menopause. Many women also face additional stresses such as responsibilities both at work and home, single parenthood, and caring for children and for aging parents.

A recent NIMH study showed that in the case of severe premenstrual syndrome (PMS), women with a preexisting vulnerability to PMS experienced relief from mood and physical symptoms when their sex hormones were suppressed. Shortly after the hormones were re-introduced, they again developed symptoms of PMS. Women without a history of PMS reported no effects of the hormonal manipulation.

Many women are also particularly vulnerable after the birth of a baby. The hormonal and physical changes, as well as the added responsibility of a new life, can be factors that lead to postpartum depression in some women. While transient "blues" are common in new mothers, a full-blown depressive episode is not a normal occurrence and requires active intervention. Treatment by a sympathetic physician and the family's emotional support for the new mother are prime considerations in aiding her to recover her physical and mental well being and her ability to care for and enjoy the infant.

Depression in Men

Although men are less likely to suffer from depression than women, three to four million men in the United States are affected by the illness. Men are less likely to admit to depression, and doctors are less likely to suspect it. The rate of suicide in men is four times that of women, though more women attempt it. In fact, after age 70, the rate of men's suicide rises, reaching a peak after age 85.

Depression can also affect the physical health in men differently from women. A new study shows that, although depression is associated with an increased risk of coronary heart disease in both men and women, only men suffer a high death rate.²

Men's depression is often masked by alcohol or drugs, or by the socially acceptable habit of working excessively long hours. Depression typically shows up in men not as feeling hopeless and helpless, but as being irritable, angry, and discouraged; hence, depression may be difficult to recognize as such in men. Even if a man realizes that he is depressed, he may be less willing than a woman to seek help. Encouragement and support from concerned family members can make a difference. In the workplace, employee assistance professionals or worksite mental health programs can be of assistance in helping men understand and accept depression as a real illness that needs treatment.

DIAGNOSTIC EVALUATION AND TREATMENT

The first step to getting appropriate treatment for depression is a physical examination by a physician. Certain medications as well as some medical conditions such as a viral infection can cause the same symptoms as depression, and the physician should rule out these possibilities through examination, interview, and lab tests. If a physical cause for the depression is ruled out, a psychological evaluation should be done, by the physician or by referral to a psychiatrist or psychologist.

A good diagnostic evaluation will include a complete history of symptoms, i.e., when they started, how long they have lasted, how severe they are, whether the patient had them before and, if so, whether the symptoms were treated and what treatment was given. The doctor should ask about alcohol and drug use, and if the patient has thoughts about death or suicide. Further, a history should include questions about whether other family members have had a depressive illness and, if treated, what treatments they may have received and which were effective.

Last, a diagnostic evaluation should include a mental status examination to determine if speech or thought patterns or memory have been affected, as sometimes happens in the case of a depressive or manic-depressive illness.

Treatment choice will depend on the outcome of the evaluation. There are a variety of antidepressant medications and psychotherapies that can be used to treat depressive disorders. Some people with milder forms may do well with psychotherapy alone. People with moderate to severe depression most often benefit from antidepressants. Most do best with combined treatment: medication to gain relatively quick symptom relief and psychotherapy to learn more effective ways to deal with life's problems, including depression. Depending on the patient's diagnosis and severity of symptoms, the therapist may prescribe medication and/or one of the several forms of psychotherapy that have proven effective for depression.

Medications

There are several types of antidepressant medications used to treat depressive disorders. These include newer medications—chiefly the selective serotonin reuptake inhibitors (SSRIs)—the tricyclics, and the monoamine oxidase inhibitors (MAOIs). The SSRIs—and other newer medications that affect neurotransmitters such as dopamine or norepinephrine—generally have fewer side effects than tricyclics. Sometimes the doctor will try a variety of antidepressants before finding the most effective medication or combination of medications. Sometimes the dosage must be increased to be effective. Although some improvements may be seen in the first few weeks, antidepressant medications must be taken regularly for 3 to 4 weeks (in some cases, as many as 8 weeks) before the full therapeutic effect occurs.

Antidepressant drugs are not habit-forming. However, as is the case with any type of medication prescribed for more than a few days, antidepressants have to be carefully monitored to see if the correct dosage is being given. The doctor will check the dosage and its effectiveness regularly.

PSYCHOTHERAPIES

Many forms of psychotherapy, including some short-term (10-20 week) therapies, can help depressed individuals. "Talking" therapies help patients gain insight into and resolve their problems through verbal exchange with the therapist, sometimes combined with "homework" assignments between sessions. "Behavioral" therapists help patients learn how to obtain more satisfaction and rewards through their own actions and how to unlearn the behavioral patterns that contribute to or result from their depression.

Two of the short-term psychotherapies that research has shown helpful for some forms of depression are interpersonal and cognitive/behavioral therapies. Interpersonal therapists focus on the patient's disturbed personal relationships that both cause and exacerbate (or increase) the depression. Cognitive/behavioral therapists help patients change the negative styles of thinking and behaving often associated with depression.

Psychodynamic therapies, which are sometimes used to treat depressed persons, focus on resolving the patient's conflicted feelings. These therapies are often reserved until the depressive symptoms are significantly improved. In general, severe depressive illnesses, particularly those that are recurrent, will require medication (or ECT under special conditions) along with, or preceding, psychotherapy for the best outcome.

HOW TO HELP YOURSELF IF YOU ARE DEPRESSED

Depressive disorders make one feel exhausted, worthless, helpless, and hopeless. Such negative thoughts and feelings make some people feel like giving up. It is important to realize that these negative views are part of the depression and typically do not accurately reflect the actual circumstances. Negative thinking fades as treatment begins to take effect. **In the meantime:**

- Set realistic goals in light of the depression and assume a reasonable amount of responsibility.
- Break large tasks into small ones, set some priorities, and do what you can as you can.
- Try to be with other people and to confide in someone; it is usually better than being alone and secretive.
- Participate in activities that may make you feel better.
- Mild exercise, going to a movie, a ballgame, or participating in religious, social, or other activities may help.
- Expect your mood to improve gradually, not immediately. Feeling better takes time.
- It is advisable to postpone important decisions until the depression has lifted. Before deciding to make a significant transition (such as) - change jobs, get married or divorced - discuss it with others who know you well and have a more objective view of your situation.
- People rarely "snap out of" a depression. But they can feel a little better day-by-day.
- Remember, positive thinking will replace the negative thinking that is part of the depression and will disappear as your depression responds to treatment.
- Let your family and friends help you.

INFORMATION

Obtained from: National Institute of Mental Health
Website: <http://www.nimh.nih.gov>